



Open Archive Toulouse Archive Ouverte

OATAO is an open access repository that collects the work of Toulouse researchers and makes it freely available over the web where possible

This is an author's version published in: <http://oatao.univ-toulouse.fr/21345>

To cite this version:

Ricci, Benoit and Lavigne, Claire and Alignier, Audrey and Aviron, Stéphanie and Biju-Duval, Luc and Bouvier, Jean-Charles and Choisis, Jean-Philippe[✉] and Franck, Pierre and Joannon, Alexandre and Ladet, Sylvie[✉] and Mezerette, Florian and Plantegenest, Manuel and Roger, Jean-Luc and Savary, Gérard and Thomas, Cecile and Vialatte, Aude[✉] and Petit, Sandrine *Efficiency of conservation biological control depends on interaction effects between landscape and pesticide use intensity.* (2018) In: *Sfécologie-2018, International Conference of Ecological Sciences*, 22 October 2018 - 25 October 2018 (Rennes, France).

Any correspondence concerning this service should be sent to the repository administrator: tech-oatao@listes-diff.inp-toulouse.fr

Efficiency of conservation biological control depends on interaction effects between landscape and pesticide use intensity

Benoit Ricci^{*1}, Claire Lavigne², Audrey Alignier³, Stéphanie Aviron³, Luc Biju-Duval¹, Jean-Charles Bouvier², Jean-Philippe Choisis⁴, Pierre Franck², Alexandre Joannon³, Sylvie Ladet⁴, Florian Mézerette¹, Manuel Plantegenest⁵, Jean-Luc Roger³, Gérard Savary³, Cécile Thomas², Aude Vialatte⁴, and Sandrine Petit¹

¹Agroécologie [Dijon] – Institut National de la Recherche Agronomique : UMRINRA 1347, Université de Bourgogne, AgroSup Dijon - Institut National Supérieur des Sciences Agronomiques, de l'Alimentation et de l'Environnement : UMR1347 – Université de Bourgogne - 17 rue Sully - BP 86510 - 21000 Dijon, France

²Unité de recherche Plantes et Systèmes de Culture Horticoles (PSH) – Institut National de la Recherche Agronomique : UR1115 – Domaine St Paul, Site Agroparc, 84941 Avignon Cedex 9, France, France

³Biodiversité agroécologie et aménagement du paysage (UMR BAGAP) – ESA - Ecole supérieure d'Agriculture d'Angers, Agrocampus Ouest, Institut National de la Recherche Agronomique – 65 rue de Saint-Brieuc 35042 Rennes, France

⁴Dynamiques Forestières dans l'Espace Rural (DYNAFOR) – Institut National Polytechnique [Toulouse], Institut National de la Recherche Agronomique : UR1201, Ecole Nationale Supérieure Agronomique de Toulouse – France

⁵Institut de Génétique, Environnement et Protection des Plantes (IGEPP) – Institut National de la Recherche Agronomique : UMR1349, Université de Rennes 1 : UMR1349, Agrocampus Ouest : UMR1349 – Domaine de la Motte au Vicomte BP 3532735653 Le Rheu, France

Abstract

Control of crop pests by their natural enemies provides a valuable but poorly quantified ecosystem service. Many studies have addressed the multiple scale drivers of pest control but results from such studies appear variable, if not equivocal. The variability of landscape responses can result from its modulation factors by local practices, among which pesticide use intensity which affects pests and natural enemies and could be a major driver of natural pest control. In this study, we examine the relative contribution of pesticide use and landscape on biological control along a double gradient of pesticide use and landscape simplification. Eighty commercial fields were monitored during three years in four contrasted regions in France. Biocontrol in each field was estimated using three different types of sentinel preys (*Ephestia kuehniella* eggs, *Acyrthosiphon pisum* aphids and *Viola arvensis* seeds). For each one, the interaction effects of landscape variables with the intensity of pesticide use on predation patterns were analyzed.

The predation of weed seeds was negatively influenced by both pesticide use and landscape

^{*}Speaker

simplification, without interaction between the two scales. The predation of *Ephestia* was negatively influenced by landscape simplification only in case of low pesticide use intensity whereas the effect was positive in case of high pesticide use intensity. There was a significant interaction between pesticide use and the length of interfaces between crops and woods on the predation aphids. These landscape variables had a positive effects on the predation of aphids in case of low or medium pesticide use intensity but a negative effect in case of high pesticide use intensity. These results could indicate that in fields with low local pesticide use, landscape complexity enhance biological control whereas in intensive fields, the presence of semi-natural habitat could be unsuitable for natural enemies adapted to intensive situations.